

Incretin Therapy and Islet Pathology: A Time for Caution

Steven E. Kahn

The past 25 years have seen the introduction of a number of new classes of medications for treating type 2 diabetes. The primary goal of these drugs is to safely lower plasma glucose in order to simultaneously reduce vascular complications and improve quality of life.

It is equally critical to demonstrate an agent's therapeutic effectiveness as it is to prove its safety. Frequently, this determination of safety occurs in the very early stages of drug development. However, in some instances, it is only after sufficient clinical experience has been obtained that the potential for harm becomes apparent. Recent examples include two therapeutically effective thiazolidinediones: troglitazone, which increased the risk of hepatic injury (1), and rosiglitazone, which a meta-analysis suggested increased the risk of cardiovascular events (2).

Two recently introduced classes of glucose-lowering agents increase incretin action. One is the glucagon-like peptide-1 (GLP-1) receptor agonists. In 2005, exenatide was the first of these agents to be introduced (3). The other is the dipeptidyl peptidase-4 (DPP-4) inhibitors, of which sitagliptin was initially approved in 2006 (3). These medications are widely used, and are associated with a reduced risk of hypoglycemia and weight gain. This, despite the fact that they may increase the risk of pancreatitis (4,5), a condition that is known to be more common in type 2 diabetes (6).

In this issue of *Diabetes*, authors from Florida and California together report pancreatic morphology from a limited series of samples from diabetic individuals who did or did not receive incretin therapy and a cohort not known to have diabetes (7). Seven of the diabetic samples were from patients who received sitagliptin and one who received exenatide, all for unknown periods of time beyond 1 year. Pancreata were procured from brain-dead organ donors by the Network for Pancreatic Organ Donors with Diabetes (nPOD), a collaborative resource funded by the Juvenile Diabetes Research Foundation to promote type 1 diabetes research (<http://www.jdrfnpod.org>).

Aside from confirming that diabetic patients who had not received incretin therapy had reduced numbers of β -cells, the study offers some new observations. First, the mass of endocrine cells was greater in subjects receiving incretin therapy. β -Cell mass was increased sixfold in diabetic

subjects receiving an incretin-based medication versus those who were not, and was threefold greater than in nondiabetic donors. Further, α -cell mass was increased fivefold in those with diabetes treated to enhance incretin action compared with those who were not, with mass in the nontreated diabetic group being comparable to those without diabetes. The increase in mass of these two endocrine cell types was due primarily to an increase in cell number as cell size was similar across groups. However, the number of β - and α -cells undergoing replication did not differ among the three groups. Second, insulin-immunoreactive cells related to ducts were present in all three groups and did not differ in frequency in the two diabetic groups. Glucagon-immunoreactive cells were present in ducts or the periductal region and formed intraductal luminal projections as observed in chronic pancreatitis. A proportion of insulin-positive cells were also positive for glucagon, the proportion being increased in both diabetic groups, but were higher in those who received sitagliptin or exenatide. Third, pancreatic mass was increased by about 40% in diabetic patients receiving incretin-based therapy. This increase in mass was accompanied by an increase in proliferation of exocrine cells as well as dysplastic changes in the form of intraepithelial neoplasia; the latter increased a little over twofold in those with diabetes who received incretin therapy versus those who did not and was frequently associated with ductal α -cell complexes. Finally, three subjects treated with sitagliptin had glucagon-producing microadenomas, and one of them also had a glucagon-producing neuroendocrine tumor. Based on these observations, the authors suggest that it is time for caution and that additional work is required to better understand these changes and perhaps learn how to harness some of them for therapeutic benefit.

Aspects of the current study and the work of others need to be considered in parallel. First, substantial differences exist between the two diabetic groups. Subjects in the control group who did not receive incretin-based therapy were 18 years younger, 67% were female, 5 were diagnosed with diabetes at the age of 20 years or younger, and 2 died of diabetic ketoacidosis. Further, five were not receiving glucose-lowering medications with the other seven on a single agent, which for four was insulin. Among those who received incretin-based therapy, only 25% were female, the group had a longer duration of diabetes, and seven of the eight subjects were using two or more medications to treat their diabetes. Thus, is the increase in pancreas mass in those receiving sitagliptin or exenatide because of these medications or is the increase more a function of some of those not receiving the medications actually having type 1 diabetes, a suggestion consistent with the greater use of insulin and much younger age at diagnosis in the latter group? This is important as magnetic resonance imaging studies have shown that pancreatic volume in subjects with type 1 diabetes is reduced by 26%

From the Division of Metabolism, Endocrinology and Nutrition, Department of Medicine, VA Puget Sound Health Care System and University of Washington, Seattle, Washington.

Corresponding author: Steven E. Kahn, skahn@u.washington.edu.

DOI: 10.2337/db13-0520

© 2013 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

See accompanying original article, p. 2595.

within months of diagnosis and by 48% after at least 10 years of the disease (8), an observation supported from weights of pancreas samples in the nPOD resource (9). Further, could the differences in body size, sex, and age be confounders that explain some of the differing morphometric observations in the three groups? A second consideration involves the observation that the profound increase in β -cell mass in those exposed to incretin-based therapies was not a result of an increase in β -cell replication. Given that the small difference in β -cell size cannot account for this change, it is not clear how this β -cell mass increase is occurring. Could it be because of the therapy reducing β -cell apoptosis (10,11), effectively negating the increased apoptosis observed in type 2 diabetes (12,13)? Such information would be valuable and would help us better understand the current findings. Or is the difference in β -cell—and possibly α -cell—mass again related to an imbalance in the types of diabetes in the two groups of diabetic subjects? Alternatively, could it be related to the preterminal clinical status of the donors, a factor that has been demonstrated to increase the rates of replication of both endocrine and nonendocrine cells in a series of 363 human organ donors on prolonged life support (14)? Third, the morphological abnormalities based on glucagon staining included microadenomas in 37.5% of the diabetic subjects treated with an incretin-based therapy and a glucagon producing neuroendocrine tumor in one of the seven who received sitagliptin. In contrast, the prevalence of pancreatic endocrine tumors is extremely low, estimated to be at 0.0005% (15). Even assuming a lower prevalence of glucagon microadenomas—say 10%—and the millions of patient-years of exposure to these agents, by now, would one not have expected reports of an increase in glucagon-related abnormalities in pancreas samples obtained at biopsy or autopsy and/or symptoms of glucagon excess beside hyperglycemia in diabetic patients on these classes of medications? Fourth, in subjects who have undergone gastric bypass surgery, postprandial GLP-1 levels are increased more than threefold (16), levels in the range of or greater than those observed with DPP-4 inhibition (17). Further, the hyperinsulinism observed in post-gastric bypass subjects

results from an increase in GLP-1-stimulated insulin secretion (18), with those experiencing hypoglycemia not having evidence of either increased β -cell mass or formation as long as 8 years after surgery (19). As many of these postsurgical subjects will have had diabetes and long-term exposure to increased levels of endogenous GLP-1, could one have expected somewhat similar findings in both endocrine and exocrine tissue in these patients to the diabetic patients in the current study? Or are the findings confounded by changes in body mass and may be an observation yet to come? Could it be—as the authors suggest—that differences in the local production of GLP-1 may exist, for which some immunostaining could have provided useful support? Fifth, while human data are always more valuable than findings in animals, we should not simply ignore substantial preclinical work that fails to substantiate a link between pancreatitis, undesired islet cell proliferation, and incretin-based therapies (20,21). This includes a consistent absence of change in the morphology and mass of α -cells in animals treated with incretin-based therapies (11,22), in contrast to marked α -cell hyperplasia and hyperglucagonemia in the mice with glucagon receptor ablation cited by the authors (23,24). Further, it is difficult to find literature substantiating the contention that partial reduction of glucagon secretion leads to compensatory α -cell hyperplasia.

The U.S. Food and Drug Administration's (FDA's) requirement that evidence of cardiovascular safety be provided for new glucose-lowering agents means that all medications developed to enhance incretin action are or will be evaluated in long-term clinical trials (Table 1). In addition, a long-term National Institutes of Health-funded study will compare two of them (sitagliptin and liraglutide) to the sulfonylurea glimepiride and insulin glargine as add-on therapy to metformin in patients with type 2 diabetes (25). A valuable byproduct of these studies will be the opportunity for adjudicating clinical events related to pancreatic pathology, be it pancreatitis or pancreatic malignancy. This approach will surely be the most informative yet, providing data obtained in a rigorous manner in patients with thousands of person-years of exposure. As none of the independent data and safety monitoring boards overseeing

TABLE 1
Long-term studies examining the safety of incretin-based therapies

Medication	Study name	ClinicalTrials.gov identifier	Comparator	Estimated number of subjects	Study start date	Estimated study end date
DPP-4 inhibitors						
Alogliptin	EXAMINE	NCT00968708	Placebo	5,400	October 2009	December 2013
Linagliptin	CAROLINA	NCT01243424	Glimepiride	6,000	October 2010	September 2018
Saxagliptin	SAVOR-TIMI53	NCT01107886	Placebo	16,500	May 2010	July 2013
Sitagliptin	TECOS	NCT00790205	Placebo	14,000	December 2008	December 2014
GLP-1 receptor agonists						
Duraglutide	REWIND	NCT01394952	Placebo	9,622	July 2011	April 2019
Exenatide	EXSCEL	NCT01144338	Placebo	9,500	June 2010	March 2017
Liraglutide	LEADER	NCT01179048	Placebo	9,340	August 2010	January 2016
Lixisenatide	ELIXA	NCT01147250	Placebo	6,000	June 2010	May 2014

Source: ClinicalTrials.gov, accessed on 27 March 2013.

the ongoing studies has terminated any of them prematurely for cause, it is doubtful they are currently observing a worrisome signal of excess pancreatic malignancy. Should there be insufficient events in each individual study, pooled data could be used for meta-analyses of these critical outcomes.

As the type 2 diabetes epidemic continues worldwide, it would seem prudent to be cautious given the findings of the current study (7). The morphological findings reported in this study should prompt the FDA and independent investigators to undertake thorough examinations of these and other pancreatic samples from patients with type 2 diabetes carefully matched for age, sex, duration of disease, and concomitant therapies who have and have not been exposed to incretin-based therapies. Further, they should reanalyze currently available data from all clinical trials with these agents. Sound clinical decision making requires the use of reproducible scientific data from well-controlled rigorous experiments that are carried out with carefully matched control groups. In this regard, the current single morphological study in a small number of poorly matched subjects is sufficient to raise important questions and prompt additional investigation. However, the current level of evidence falls short of that required to prematurely banish two novel therapeutic classes that have thus far proven to be valuable in treating type 2 diabetes.

ACKNOWLEDGMENTS

This work was supported in part by the U.S. Department of Veterans Affairs and National Institutes of Health Grant DK-017047.

S.E.K. has received fees for consulting on incretin-based therapy from Boehringer Ingelheim, Bristol-Myers Squibb, Eli Lilly, GlaxoSmithKline, Intarcia, Merck, and Novo Nordisk. No other potential conflicts of interest relevant to this article were reported.

REFERENCES

- Misbin RI. Troglitazone-associated hepatic failure. *Ann Intern Med* 1999; 130:330
- Nissen SE, Wolski K. Effect of rosiglitazone on the risk of myocardial infarction and death from cardiovascular causes. *N Engl J Med* 2007; 356:2457–2471
- Drucker DJ, Nauck MA. The incretin system: glucagon-like peptide-1 receptor agonists and dipeptidyl peptidase-4 inhibitors in type 2 diabetes. *Lancet* 2006;368:1696–1705
- Elashoff M, Matveyenko AV, Gier B, Elashoff R, Butler PC. Pancreatitis, pancreatic, and thyroid cancer with glucagon-like peptide-1-based therapies. *Gastroenterology* 2011;141:150–156
- Singh S, Chang HY, Richards TM, Weiner JP, Clark JM, Segal JB. Glucagonlike peptide 1-based therapies and risk of hospitalization for acute pancreatitis in type 2 diabetes mellitus: a population-based matched case-control study. *JAMA Intern Med* 2013;173:534–539
- Noel RA, Braun DK, Patterson RE, Bloomgren GL. Increased risk of acute pancreatitis and biliary disease observed in patients with type 2 diabetes: a retrospective cohort study. *Diabetes Care* 2009;32:834–838
- Butler AE, Campbell-Thompson M, Gurlo T, Dawson DW, Atkinson M, Butler PC. Marked expansion of exocrine and endocrine pancreas with incretin therapy in humans with increased exocrine pancreas dysplasia and the potential for glucagon-producing neuroendocrine tumors. *Diabetes* 2013;62:2595–2604
- Williams AJ, Thrower SL, Sequeiros IM, et al. Pancreatic volume is reduced in adult patients with recently diagnosed type 1 diabetes. *J Clin Endocrinol Metab* 2012;97:E2109–E2113
- Campbell-Thompson M, Wasserfall C, Montgomery EL, Atkinson MA, Kaddis JS. Pancreas organ weight in individuals with disease-associated autoantibodies at risk for type 1 diabetes. *JAMA* 2012;308:2337–2339
- Aston-Mourney K, Hull RL, Zraika S, Udayasankar J, Subramanian SL, Kahn SE. Exendin-4 increases islet amyloid deposition but offsets the resultant beta cell toxicity in human islet amyloid polypeptide transgenic mouse islets. *Diabetologia* 2011;54:1756–1765
- Takeda Y, Fujita Y, Honjo J, et al. Reduction of both beta cell death and alpha cell proliferation by dipeptidyl peptidase-4 inhibition in a streptozotocin-induced model of diabetes in mice. *Diabetologia* 2012;55:404–412
- Butler AE, Janson J, Bonner-Weir S, Ritzel R, Rizza RA, Butler PC. β -Cell deficit and increased β -cell apoptosis in humans with type 2 diabetes. *Diabetes* 2003;52:102–110
- Jurgens CA, Toukatly MN, Fligner CL, et al. β -Cell loss and β -cell apoptosis in human type 2 diabetes are related to islet amyloid deposition. *Am J Pathol* 2011;178:2632–2640
- Int' Veld P, De Munck N, Van Belle K, et al. β -Cell replication is increased in donor organs from young patients after prolonged life support. *Diabetes* 2010; 59:1702–1708
- Oberg K. Pancreatic endocrine tumors. *Semin Oncol* 2010;37:594–618
- Korner J, Bessler M, Inabnet W, Taveras C, Holst JJ. Exaggerated glucagon-like peptide-1 and blunted glucose-dependent insulinotropic peptide secretion are associated with Roux-en-Y gastric bypass but not adjustable gastric banding. *Surg Obes Relat Dis* 2007;3:597–601
- Herman GA, Bergman A, Stevens C, et al. Effect of single oral doses of sitagliptin, a dipeptidyl peptidase-4 inhibitor, on incretin and plasma glucose levels after an oral glucose tolerance test in patients with type 2 diabetes. *J Clin Endocrinol Metab* 2006;91:4612–4619
- Salehi M, Prigeon RL, D'Alessio DA. Gastric bypass surgery enhances glucagon-like peptide 1-stimulated postprandial insulin secretion in humans. *Diabetes* 2011;60:2308–2314
- Meier JJ, Butler AE, Galasso R, Butler PC. Hyperinsulinemic hypoglycemia after gastric bypass surgery is not accompanied by islet hyperplasia or increased beta-cell turnover. *Diabetes Care* 2006;29:1554–1559
- Koehler JA, Baggio LL, Lamont BJ, Ali S, Drucker DJ. Glucagon-like peptide-1 receptor activation modulates pancreatitis-associated gene expression but does not modify the susceptibility to experimental pancreatitis in mice. *Diabetes* 2009;58:2148–2161
- Nyborg NC, Mølck AM, Madsen LW, Knudsen LB. The human GLP-1 analog liraglutide and the pancreas: evidence for the absence of structural pancreatic changes in three species. *Diabetes* 2012;61:1243–1249
- Uhles S, Wang H, Bénardeau A, et al. Taspoglutide, a novel human once-weekly GLP-1 analogue, protects pancreatic β -cells in vitro and preserves islet structure and function in the Zucker diabetic fatty rat in vivo. *Diabetes Obes Metab* 2011;13:326–336
- Gelling RW, Du XQ, Dichmann DS, et al. Lower blood glucose, hyperglucagonemia, and pancreatic alpha cell hyperplasia in glucagon receptor knockout mice. *Proc Natl Acad Sci USA* 2003;100:1438–1443
- Yu R, Dhall D, Nissen NN, Zhou C, Ren SG. Pancreatic neuroendocrine tumors in glucagon receptor-deficient mice. *PLoS ONE* 2011;6:e23397
- Nathan DM, Buse JB, Kahn SE, et al; GRADE Study Research Group. Rationale and design of the Glycemia Reduction Approaches in Diabetes: A Comparative Effectiveness Study (GRADE). *Diabetes Care*. 20 May 2013 [Epub ahead of print]